

ASYMPTOTICS OF ORTHOGONAL POLYNOMIALS BEYOND THE SCOPE OF SZEGŐ'S THEOREM

F. PEHERSTORFER, A. VOLBERG, P. YUDITSKII

ABSTRACT. First we give here a simple proof of a remarkable result of Videnskii and Shirokov: let B be a Blaschke product with n zeros, then there exists an outer function $\phi, \phi(0) = 1$, such that $\|(B\phi)'\| \leq Cn$, where C is an absolute constant. Then we apply this result to a certain problem of finding the asymptotic of orthogonal polynomials.

1. INTRODUCTION AND THE MAIN RESULTS

Let us introduce the quantity investigated in Videnskii-Shirokov paper [6].

Let X be a Banach space of functions analytic in the open unit disk $\mathbb{D}$, let E be a finite collection of points in $\mathbb{D}$, and let B_E denote the Blaschke product vanishing precisely at the points of E . Characterization of zero sets of functions from X is a difficult and important problem. It is closely related to the estimation of the quantity

$$\phi(E, X) := \inf\{\|f\| : f \in X, f = B_E G, G(0) = 1\}.$$

The following sequence plays an important part in understanding zero sets structure of X :

$$r_n(X) := \sup\{\phi(E, X) : \text{card} E = n\}.$$

Very interesting research is related to asymptotics of $r_n(l_a^1)$, where l_a^1 stands for analytic functions with absolutely summable Taylor coefficients. In the papers of Shäffer [5], Gluskin, Meyer, Pajor [1] and Queffelec [4] it was proved that $r_n(l_a^1) \asymp \sqrt{n}$. Actually Shäffer, motivated by Van der Waerden, studied the asymptotics of another quantity, which turns out to be equal to $r_n(l_a^1)$ (see [5], [1], [2]). In [2] one can find an easy proof of equality of $r_n(l_a^1)$ and this other quantity

$$k_n := \sup\{|\det T| \|T^{-1}\| : \|T\| \leq 1\},$$

where the supremum is taken over all invertible matrices of order n and over *all norms*! But if one fixes the norm of T as acting in n -dimensional

Partially supported by NSF grant DMS-0200713.

AMS subject classification codes: 42B20, 42C15.

Key words: Orthogonal polynomials, extremal problems, Blaschke product, CMV matrices.

dimensional Banach space one can consider the analog of k_n , or the following problem of finding the best estimate

$$\|T^{-1}\| \leq \Phi_n(\delta),$$

where we consider all invertible T in n -dimensional Banach space A , such that $\|T\| \leq 1$ and the spectral radius $r(T^{-1}) = \delta$. Those estimates are considered in details in the paper of Nikolski [2]. In [2] the following variant of the problem is also considered: the requirement $\|T\| \leq 1$ is replaced by

$$\|f(T)\| \leq C\|f\|,$$

for every polynomial f . In other words operator T is assumed to satisfy A -functional calculus with respect to function space (algebra) A . It is easy to see that for Banach algebras $A = X$ this brings us to estimates of yet another interesting quantity:

$$\text{cap}_X(E) := \inf\{\|f\|_X : f(0) = 1, f|_E = 0\}.$$

It is immediate that our quantities $\phi(E, X)$, $\text{cap}_X(E)$ are essentially related to each other.

In [2] the sharp estimates of $\text{cap}_{B_{p,q}^s}(E)$ is given. Here $B_{p,q}^s$ is a Besov class (see definition below). Notice that for non-integer s

$$B_{\infty,\infty}^s = \Lambda^s,$$

but for integer s only the strict inclusion holds:

$$(1) \quad \Lambda^s \subset B_{\infty,\infty}^s.$$

One of the results of [2] formulated in the form convenient for us is

$$r_n(B_{\infty,\infty}^s) \asymp n^s.$$

By our previous remark this gives $r_n(\Lambda^s) \asymp n^s$ for non-integer s . The question for integer s seems to be more subtle. In fact, the proof of this fact for integers given in the paper of Videnskii and Shirokov ([6], Theorem 3.1) is quite involved.

We want to give here a very simple proof. It follows the ideas of [2], [6], but it is quite short. We treat the case $s = 1$.

Theorem 1.1. *Let B be a Blaschke product with n zeros in the disc $\mathbb{D}$. Then there exists an outer function ϕ satisfying the following properties:*

$$(2) \quad \|(B\phi)'\|_{L^\infty(\mathbb{T})} \leq Cn.$$

$$(3) \quad \phi(0) = 1.$$

$$(4) \quad \|\phi\|_{L^\infty(\mathbb{T})} \leq A.$$

Here C, A are absolute constants.

1.1. **Classes** $B_{p,q}^s, \Lambda^s$. Let W_n are Valee-Poussin kernels defined by their Fourier coefficients as follows. $W_0(z) = 1 + z$, and for $n \geq 1$ $\hat{W}_n = 1$ for $k = 2^n$, $\hat{W}_n = 0$ for $k \notin (2^{n-1}, 2^{n+1})$, and $\hat{W}_n$ is affine on $[2^{n-1}, 2^n]$ and on $[2^n, 2^{n+1}]$.

Now the class $B_{p,q}^s$ consists of analytic in $\mathbb{D}$ functions $f(z) = \sum \hat{f}(n)z^n$ such that the sequence $\{2^{ns}\|f \star W_n\|_{L^p(\mathbb{T})}\}$ belongs to l^q .

For integer $s = n$ the class Λ^s consists of all functions analytic in $\mathbb{D}$ for which the following seminorm is finite ??

$$\|f\|_{\Lambda^n} := \|f^{(n)}\|_{H^\infty}.$$

If $s = n + \alpha$, $0 < \alpha < 1$, then $f \in \Lambda^s$ means that the previous seminorm is finite and in addition to that one more seminorm should be finite:

$$\|f^{(n)}\|_{\Lambda^\alpha} := \sup_{z, \zeta \in \mathbb{D}, z \neq \zeta} \frac{|f^{(n)}(z) - f^{(n)}(\zeta)|}{|z - \zeta|^\alpha}.$$

2. PROOF OF THEOREM 1.1

Proof. Let $\tilde{B}$ be a Blaschke product with the same zeros $\{z_1, \dots, z_n\}$ as B but in $(1 + \frac{1}{n})\mathbb{D}$. If $R = 1 + \frac{1}{n}$ then

$$\tilde{B} = \prod_{k=1}^n \frac{\frac{z}{R} - \frac{\bar{z}_k}{R}}{1 - \frac{1}{R^2} \bar{z}_k z}.$$

We can write

$$R^n \tilde{B} =: B\phi_0,$$

having this as a definition of ϕ_0 .

So

$$\phi_0(z) = \prod_{k=1}^n \frac{1 - \bar{z}_k z}{1 - \frac{1}{R^2} \bar{z}_k z}.$$

Obviously, if $z \in \mathbb{T}$ then $|\tilde{B}| < 1$, and, therefore,

$$|\phi_0(z)| \leq (1 + \frac{1}{n})^n \leq A.$$

So (4) is satisfied. Also $\phi_0(0) = 1$. To check (2) we write the Cauchy formula for $\tilde{B}(\zeta_0), \zeta_0 \in \mathbb{T}$:

$$\tilde{B}(\zeta_0) = \frac{1}{2\pi i} \int_{(1+\frac{1}{n})\mathbb{T}} \frac{\tilde{B}(\zeta)}{\zeta - \zeta_0} d\zeta.$$

Hence

$$|\tilde{B}'(\zeta_0)| \leq \frac{1}{2\pi} \int_{(1+\frac{1}{n})\mathbb{T}} \frac{|\tilde{B}(\zeta)|}{|\zeta - \zeta_0|^2} dm(\zeta).$$

And therefore

$$|\tilde{B}'(\zeta_0)| \leq C \int_{(1+\frac{1}{n})\mathbb{T}} \frac{1}{|\zeta - \zeta_0|^2} dm(\zeta) \leq C n.$$

But then

$$|(B\phi_0)'(\zeta_0)| = (1 + \frac{1}{n})^n |\tilde{B}'(\zeta_0)| \leq AC n.$$

□

Remarks. 1. Replacing $R = 1 + \frac{1}{n}$ by $R = 1 + \frac{\varepsilon}{n}$ we can improve the estimate of ϕ_0 : $\|\phi_0\|_\infty \leq 1 + \varepsilon$ by paying the price in (2), the constant begins to depend on ε :

$$(5) \quad \|(B\phi)'\|_{L^\infty(\mathbb{T})} \leq C_\varepsilon n.$$

2. We proved also that

$$\|B\phi_0\|_{\Lambda^s} \leq C n^s.$$

For $s \in (0, 1)$ this follows immediately from (2) and (4). For $s = 1$ we just presented the proof. For $s > 1$ the proof goes exactly along the same lines but we need to differentiate more times.

The second remark contains the claim of Theorem 3.1 of [6].

The first remark plays a crucial part in this paper, where we consider the asymptotics of polynomials orthogonal with respect to measure μ of the following type

$$d\mu := \frac{dm}{|\psi(e^{i\theta})|^2} + \sum_{k=1}^{\infty} \mu_k \delta_{z_k},$$

where z_k form the sequence of Blaschke.

In this situation one should distinguish between the sequence of Blaschke inside $\mathbb{D}$ and outside $\mathbb{D}$. The second case is much more interesting as expected asymptotics of $L^2(\mu)$ -orthogonal polynomials is different from the classical asymptotics of Szegő, where all $\mu_k = 0$.

The orthogonalization with respect to measures described above is an attempt to generalize the results of the paper of Peherstorfer and Yuditskii [3], where the "point spectrum" was assumed to be lying on the real line. This is quite an interesting paper which extends the strong asymptotic results for orthogonal polynomials that satisfy a Szegő condition on a real interval to the case where the measure has in addition a denumerable set of mass points outside the interval. The authors assume that the mass points outside of the interval accumulate only at the endpoints of the interval and that a Blaschke condition is satisfied. Under these conditions, strong asymptotic results are given for the orthonormal polynomials and for their leading coefficients. The asymptotic behavior is stated in terms of the Szegő function for the absolutely continuous part and of a Blaschke product related to the extra mass points.

3. ORTHOGONAL POLYNOMIALS AND SMOOTHNESS OF $B\phi$

Let μ be as above with

$$|z_k| > 1, \sum_k (|z_k| - 1) < \infty,$$

and let $\{P_n\}$ denote the sequence of analytic polynomials orthonormal with respect to μ .

We are writing

$$P_n(z) = \tau_n z^n + \dots + a_{0n}$$

and we want to find the limit of τ_n (we will prove that it exists in many cases).

We also consider rational functions of the form

$$(6) \quad R_{n,m}(z) = \eta_{n,m} z^n + \dots + a_{0n} + \dots + a_{m,n} z^m,$$

where $m = -n$ or $-(n-1)$. We want them to be mutually orthogonal in $L^2(\mu)$ and of norm 1 in this space. We wish to discuss the asymptotics of the “leading” coefficient $\eta_n := \eta_{n, -(n-1)}$.

Here are our two main results. We have the convention that $\tau_n, \eta_n = \eta_{n, -(n-1)}$ are all positive, and that $B(0), \psi(0)$ are also positive. The first theorem deals with the perturbation of so-called CMV matrices.

Theorem 3.1. *For any $\{z_k\}$ satisfying the Blaschke condition as above, for any summable $\{\mu_k\}$, and for ψ bounded away from 0 one has*

$$\lim_{n \rightarrow \infty} \eta_n = B(0)\psi(0).$$

The asymptotics of τ_n seems to be more subtle. For certain geometric configuration of $\{z_k\}$ no extra assumptions on $\{\mu_k\}$ is needed. For example, if $\{z_k\}$ converge in a Stolz star to only a closed set $E \subset \mathbb{T}$ satisfying $m(E) = 0$ and $\sum l_n \log \frac{1}{l_n} < \infty$ (here l_n stand for lengths of complimentary intervals of E ; such sets are called Carleson subsets of $\mathbb{T}$) we do not need any extra assumption on $\{\mu_k\}$. A particular case of such a geometry is the case of real $\{z_k\}$ considered in [3]. But if $\{z_k\}$ is an arbitrary Blaschke sequence outside of $\mathbb{D}$, we still need one assumption on $\{\mu_k\}$ saying that the series $\sum_k \mu_k$ converges with a definite speed. Here is this assumption:

$$(7) \quad \sum_{1 < |z_k| < 1 + \frac{1}{n}} \mu_k \leq C_A \frac{1}{(\log n)^A}, \quad \forall A.$$

Theorem 3.2. *For any $\{z_k\}$ satisfying the Blaschke condition as above, for any summable $\{\mu_k\}$ satisfying the extra condition (7), and for ψ bounded away from 0 and from ∞ one has*

$$\lim_{n \rightarrow \infty} \eta_n = B(0)\psi(0).$$

4. THE PROOF OF THEOREM 3.1

Lemma 4.1. *Let R_n be a rational function solving the following extremal problem: find*

$$\sup\{|\eta| : \mathcal{R}(z) = \eta z^n + \dots + az^{-(n-1)}, \|R\|_{L^2(\mu)} \leq 1\}.$$

Then $R_n := R_{n, -(n-1)}$ from (6) solves this extremal problem.

Proof. Consider another extremal problem: $\inf\{\|Q\|_{L^2(\mu)} : \mathcal{Q}(z) = z^n + \dots + az^{-(n-1)}\}$. These are $L^2(\mu)$ orthogonal functions of course. Our extremal functions $\mathcal{R}$ are just $\mathcal{Q}/\|\mathcal{Q}\|_{L^2(\mu)}$, so $\inf\|\mathcal{Q}\|_{L^2(\mu)}$ means $\sup|\tau|$. $\square$

4.1. Estimate from below of η_n . Let $\zeta_k = 1/\bar{z}_k$, B be a Blaschke product with zeros $\{\zeta_k\}_{k=1}^\infty$, B_n be a partial Blaschke product built by

$$\zeta_k : |\zeta_k| < 1 - \frac{1}{k_n},$$

where $k_n := [\varepsilon_n n]$, $\varepsilon \rightarrow 0$.

We choose ε_n will be chosen later.

Without the loss of generality we think that

$$\sum_k (1 - |\zeta_k|) < 1.$$

The number of zeros of B_n is at most k_n .

Choose $k_n := [A_n \varepsilon_n n]$, where A_n grows to infinity very slow, namely, $A_n \varepsilon \rightarrow 0$. Put

$$R := 1 + \frac{1}{l_n}.$$

Let $\tilde{B}$ be a Blaschke product with the same zeros as B_n , but a Blaschke product in $R\mathbb{D}$. Then we introduce ϕ by formula (as in Theorem 1.1)

$$(8) \quad R^{k_n} \tilde{B} =: B\phi,$$

or

$$\phi := \prod_{k=1}^{k_n} \frac{1 - \bar{\zeta}_k z}{1 - \frac{1}{R^2} \bar{\zeta}_k z}.$$

In Theorem 1.1 we proved that

$$\|B\phi\|_{\Lambda^1} \leq Cl_n.$$

Therefore,

$$\|B\phi\|_{B_{\infty, \infty}^1} \leq Cl_n.$$

Let 2^k be the largest such number smaller than n . Consider $W_k, W_{k+1}, \dots$ (we introduced Vallée-Poussin kernels in Section 1). We just noticed that

$$(9) \quad \|W_j \star B\phi\|_{\infty} \leq Cl_n 2^{-j}, \quad j \geq k.$$

So (9) gives

$$(10) \quad \left\| \sum_{j \geq k} W_j \star B\phi \right\|_{\infty} \leq C l_n / n$$

And, therefore, if V_k denote the modified Vallée-Poussin kernel

$$\hat{V}_k(j) = 1, j \in [-2^{k-1}, 2^{k-1}], \quad \hat{V}_k(j) = 1, |j| \geq 2^k,$$

and $\hat{V}_k(j)$ is affine otherwise, then from (10) we conclude

$$(11) \quad \|V_k \star (B\phi) - B\phi\|_{\infty} \leq C l_n / n$$

Let us consider the following modified Vallée-Poussin kernel

$$\hat{V}P_n(j) = 1, j \in [-n, n], \quad \hat{V}P_n(j) = 1, |j| \geq n,$$

and $\hat{V}P_n(j)$ is affine otherwise. It is easy to see that we proved (using $V_k \star V P_n = V_k$)

$$(12) \quad \|V P_n \star (B\phi) - B\phi\|_{\infty} \leq 4C l_n / n \leq C' \delta_n,$$

where $\delta_n \rightarrow 0$.

Notice the estimate from above for ϕ (keeping in mind $l_n \approx A_n k_n$):

$$(13) \quad |(B_n \phi)(z)| = (1 + \frac{1}{l_n})^{k_n} |\tilde{B}(\zeta)| \leq e^{C/A_n} =: 1 + \delta'_n, \quad \delta'_n \rightarrow 0.$$

Consider $G_n = V P_n \star (B\phi)$, $r_n = G_n / z^n$. Clearly (using $\phi(0) = 1$)

$$(14) \quad |G_n(0) - B(0)\phi(0)| = |G_n(0) - B(0)| \leq \delta_n.$$

The rational function r_n is of the type we want, but only after the application of the symmetry $n \rightarrow -n$ on $\mathbb{Z}$.

Let us estimate the norm

$$\|r_n\|^2 := \int_{\mathbb{T}} |r_n|^2 dm + \sum_{k=1}^{\infty} \mu_k |r_n(\zeta_k)|^2.$$

From (12) and (13) we have $|r_n| \leq 1 + C(\delta_n + \delta'_n)$ on $\mathbb{T}$. So $\int_{\mathbb{T}} |r_n|^2 dm \leq 1 + C\delta''_n$, $\delta''_n \rightarrow 0$. Now we split the sum to two: $\alpha_1 := \sum_{k=1}^{k_n} \mu_k |r_n(\zeta_k)|^2$, $\alpha_2 := \sum_{k > k_n} \mu_k |r_n(\zeta_k)|^2$.

To estimate α_1 notice that (12) says that $G_n - B\phi$ is small on $\mathbb{T}$, and the construction of G_n says that $(G_n - B\phi)(z)$ has zero of multiplicity n at the origin. This is analytic function in $\mathbb{D}$, so by classical Schwartz lemma

$$(15) \quad |G_n(z) - B\phi(z)| \leq C\delta_n |z|^n,$$

where $\delta_n \rightarrow 0$.

In particular,

$$(16) \quad |(V P_n \star B\phi)(\zeta_k) - B\phi(\zeta_k)| \leq C\delta_n |\zeta_k|^n.$$

But $B(\zeta_k) = 0$, $k = 1, \dots, k_n$, so we rewrite

$$(17) \quad |G_n(\zeta_k)| = |(VP_n \star B\phi)(\zeta_k)| \leq C\delta_n |\zeta_k|^n, \quad k = 1, \dots, k_n.$$

And finally, using $r_n(z) = G_n(z)/z^n$, we get

$$(18) \quad |r_n(\zeta_k)| \leq C\delta_n, \quad k = 1, \dots, k_n.$$

Therefore,

$$(19) \quad |\alpha_1| \leq C\delta_n \sum_k \mu_k \leq C\delta_n.$$

To estimate α_2 we start from (16) to write

$$(20) \quad |r_n(\zeta_k)| \leq C\delta_n + |(B\phi)(\zeta_k)|/|\zeta_k|^n, \quad k > k_n.$$

But if $k > k_n$ then $1 - |\zeta_k| < \frac{C}{\varepsilon_n n}$, so

$$\frac{1}{|\zeta_k|^n} \leq e^{C/\varepsilon_n}.$$

In particular,

$$|\alpha_2| \leq \sum_{k > k_n} \mu_k |r_n(\zeta_k)|^2 \leq C\delta_n + e^{C/\varepsilon_n} \sum_{k > k_n} \mu_k.$$

We can always choose ε_n to decrease so slowly that $\sum_{k > k_n} \mu_k$ kills the growth of e^{C/ε_n} and the last expression

$$\gamma_n := e^{C/\varepsilon_n} \sum_{k > k_n} \mu_k$$

tends to zero.

Summarizing all that

$$(21) \quad \|r_n\| \leq 1 + \text{small}$$

and

$$(22) \quad r_n(z) = \rho z^{-n} + \dots + a z^{n-1}, \quad |\rho| \geq B(0) - \text{small}.$$

Let us inverse the variable. Consider $R(z) := r_n^*(z) := \overline{r(1/\bar{z})} = \bar{\rho} z^n + \dots + \bar{a} z^{-(n-1)}$. Then we just got $|\rho| \geq B(0)\phi(0) - \text{small}$ and

$$(23) \quad \|R\|_{L^2(\mu)} \leq 1 + \text{small}$$

We finished the estimate

$$(24) \quad \liminf_{n \rightarrow \infty} |\eta_n| \geq B(0).$$

This proves the estimate from below for η_n for the case $\psi = 1$. For non-trivial ψ bounded away from zero, we approximate it from below by smooth ψ' and repeat the above argument for $B\phi/\psi$ instead of $B\phi$.

5. ESTIMATE FROM BELOW OF τ_n

We repeat the beginning of Section 4.1. Let $\zeta_k = 1/\bar{z}_k$, B be a Blaschke product with zeros $\{\zeta_k\}_{k=1}^\infty$, B_n be a partial Blaschke product built by

$$\zeta_k : |\zeta_k| < 1 - \frac{1}{k_n},$$

where $k_n := [\varepsilon_n n]$, $\varepsilon \rightarrow 0$.

We choose ε_n later.

Without the loss of generality we think that

$$\sum_k (1 - |\zeta_k|) < 1.$$

The number of zeros of B_n is at most k_n .

Choose $k_n := [A_n \varepsilon_n n]$, where A_n grows to infinity very slow, namely, $A_n \varepsilon \rightarrow 0$. Put

$$R := 1 + \frac{1}{l_n}.$$

Let $\tilde{B}$ be a Blaschke product with the same zeros as B_n , but a Blaschke product in $R\mathbb{D}$. Then we introduce ϕ by formula (as in Theorem 1.1)

$$(25) \quad R^{k_n} \tilde{B} =: B\phi,$$

or

$$\phi := \prod_{k=1}^{k_n} \frac{1 - \bar{\zeta}_k z}{1 - \frac{1}{R^2} \bar{\zeta}_k z}.$$

We use all the notations of the previous section. Let T_n denote the Taylor polynomial of degree n of $B_n\phi$. Our goal is very simple, we want to repeat all the reasoning of the previous section, but instead of estimating $|G_n(z) - B\phi(z)| = |(VP_n \star B\phi)(z) - B\phi(z)|$ we wish to estimate $|T_n(z) - B\phi(z)| = |(D_n \star B\phi)(z) - B\phi(z)|$. Here D_n stands for Dirichlet kernel. In other words we just want to replace VP_n kernel by D_n kernel.

Let $\zeta_0 \in \mathbb{T}$. One can write the Cauchy formula for $B\phi(\zeta_0)$ over $R\mathbb{T}$, $R = 1 + \frac{1}{l_n}$:

$$(B_n\phi - T_n)(\zeta_0) = \frac{1}{2\pi i} \int_{(1+\frac{1}{l_n})\mathbb{T}} \frac{\zeta_0^n (B_n\phi)(\zeta)}{(\zeta - \zeta_0)\zeta^n} d\zeta.$$

We saw in (13) that

$$(26) \quad |B_n\phi(\zeta)| = (1 + \frac{1}{l_n})^{k_n} |\tilde{B}(z)| \leq e^{C/A_n} =: 1 + \kappa_n, \kappa_n \rightarrow 0.$$

In particular, putting this in our Cauchy formula we get

$$\|B_n\phi - T_n\|_\infty \leq \frac{C}{(1 + \frac{1}{l_n})^n} \int_{(1+\frac{1}{l_n})\mathbb{T}} \frac{|d\zeta|}{|\zeta - \zeta_0|},$$

or

$$\|B_n\phi - T_n\|_\infty \leq C \log l_n e^{-\frac{1}{A_n \varepsilon_n}} \leq C' \log n e^{-\frac{1}{A_n \varepsilon_n}}.$$

We denote $\delta_n := \log n e^{-\frac{1}{A_n \varepsilon_n}}$. Unlike δ_n from the previous section, this one should not go to zero. But we require that it does go to zero, thus putting some restriction on ε_n, A_n . So from now on we *assume*

$$(27) \quad \delta_n := \log n e^{-\frac{1}{A_n \varepsilon_n} \rightarrow 0},$$

and

$$(28) \quad \|B_n \phi - T_n\|_\infty \leq C \delta_n \rightarrow 0.$$

Also introduce

$$p_n(z) := T_n(z)/z^n, \quad P_n(z) := \overline{p_n(\frac{1}{\bar{z}})} = z^n \overline{T(\frac{1}{\bar{z}})}.$$

Then, let us consider the norm

$$\|p_n\| := \int_{\mathbb{T}} |p_n|^2 dm + \sum_k \mu_k |p_n(\zeta_k)|^2.$$

Of course we want to prove

$$(29) \quad \|p_n\| \leq 1 + \text{small}$$

and

$$(30) \quad |T_n(0)| \geq B(0) - \text{small}$$

Relation (30) follows from (28) and $\phi(0) = 1$.

To prove (29) we notice that (26) implies

$$\int_{\mathbb{T}} |p_n|^2 dm \leq 1 + C(\kappa_n + \delta_n) \leq 1 + \text{small}.$$

We split the sum $\sum_k \mu_k |p_n(\zeta_k)|^2$ into two: $\beta_1 := \sum_{k=1}^{k_n} \mu_k |p_n(\zeta_k)|^2$, $\beta_2 := \sum_{k > k_n} \mu_k |p_n(\zeta_k)|^2$.

To estimate β_1 we notice (28) and the fact that $B\phi - T_n$ is an analytic function in $\mathbb{D}$ with zero of multiplicity n at the origin. So by Schwartz lemma

$$(31) \quad |T_n(z) - B\phi(z)| \leq C \delta_n |z|^n,$$

where $\delta_n \rightarrow 0$.

In particular,

$$(32) \quad |T_n(\zeta_k) - B\phi(\zeta_k)| \leq C \delta_n |\zeta_k|^n.$$

But $B(\zeta_k) = 0$, $k = 1, \dots, k_n$, so we rewrite

$$(33) \quad |T_n(\zeta_k)| \leq C \delta_n |\zeta_k|^n, \quad k = 1, \dots, k_n.$$

And finally, using $p_n(z) = T_n(z)/z^n$, we get

$$(34) \quad |p_n(\zeta_k)| \leq C \delta_n, \quad k = 1, \dots, k_n.$$

Therefore,

$$(35) \quad |\alpha_1| \leq C\delta_n \sum_k \mu_k \leq C\delta_n.$$

To estimate β_2 we start from (32) to write

$$(36) \quad |p_n(\zeta_k)| \leq C\delta_n + |(B\phi)(\zeta_k)|/|\zeta_k|^n, \quad k > k_n.$$

But if $k > k_n$ then $1 - |\zeta_k| < \frac{C}{\varepsilon_n n}$, so

$$\frac{1}{|\zeta_k|^n} \leq e^{C/\varepsilon_n}.$$

In particular,

$$|\beta_2| \leq \sum_{k > k_n} \mu_k |p_n(\zeta_k)|^2 \leq C\delta_n + e^{C/\varepsilon_n} \sum_{k > k_n} \mu_k.$$

We cannot always choose ε_n to decrease so slowly that $\sum_{k > k_n} \mu_k$ kills the growth of e^{C/ε_n} and the last expression

$$(37) \quad \gamma_n := e^{C/\varepsilon_n} \sum_{k > k_n} \mu_k$$

tends to zero. The reason is in the fact that we have a restriction (27) that makes ε_n to tend to zero with a *certain* speed. But combining (27), (37), we see that they are reconcilable if μ_k tend to zero sufficiently rapidly.

It is easy to deduce that (7) allows us to have both assumptions (27), (37). Therefore, we get (29) and (30). Inversing the variable, that is passing from $p_n(z)$ to $P_n(z)$ proves

$$\liminf_{n \rightarrow \infty} |\tau_n| \geq B(0).$$

If ψ is bounded away from zero we approximate it from below by a smooth function, and thinking now that $1/\psi$ is smooth, we can repeat our approximation arguments with $B\phi/\psi$ instead of $B\phi$.

6. ESTIMATE FROM ABOVE OF τ_n AND η_n

This is a simple estimate. For η_n it is still slightly more difficult, so we show it in this case.

We want to prove now

$$(38) \quad \limsup_{n \rightarrow \infty} |\tau_n| \leq B(0)\psi(0).$$

Let $\psi_*(z) := \overline{\psi(1/\bar{z})}$, $B_*(z) := \overline{B(1/\bar{z})}$. Let B_*^k be a finite Blaschke product that is subproduct of B_* .

Consider

$$\int_{\mathbb{T}} \frac{R_n(z)}{\psi_*(z)} \overline{z^n B_*^k(z)} dm(z),$$

which absolute value evidently does not exceed one. Let us calculate the integral according to Cauchy theorem in the exterior of $\mathbb{D}$. We can do that

as R_n/z^{n+1} is analytic in the exterior of the unit disc and equals zero at infinity.

$$\int_{\mathbb{T}} \frac{R_n(z)}{\psi_*(z)z^{n+1}B_*^k(z)} dz = \frac{\eta_n}{B(0)\psi(0)} - \sum \frac{R_n(z_i)}{(B_*^k)'(z_i)\psi_*(z_i)z_i^{n+1}}.$$

The last sum we have the estimate (since R_n is normalized in $L^2(\mu)$):

$$\left| \sum \frac{R_n(z_i)}{(B_*^k)'(z_i)\psi_*(z_i)z_i^{n+1}} \right|^2 \leq \sum \frac{1}{|(B_*^k)'(z_i)\psi_*(z_i)z_i^{n+1}|^2 \mu_i}.$$

The number of terms in the sum is finite, so the sum goes to zero when n tends to infinity.

Thus,

$$\limsup_{n \rightarrow \infty} |\tau_n| \leq B_*^k(0)\psi(0).$$

Since B_*^k is arbitrary we proved (38).

Of course we the reader can see that the proof for τ_n is exactly the same.

Our both theorems are completely proved.

REFERENCES

- [1] E. Gluskin, M. Meyer, A. Pajor, Zeros of analytic functions and norms of inverse matrices, *Israel J. Math.* **87** (1994), pp. 225–242.
- [2] N. K. Nikolski, Condition numbers of large matrices and analytic capacities, Preprint, Univ. Bordeaux I, 2005.
- [3] F. Peherstorfer, P. Yuditskii *Asymptotics of orthonormal polynomials in the presence of a denumerable set of mass points*, *Proc. Amer. Math. Soc.* **129** (2001), no. 11, 3213–3220.
- [4] H. Queffelec, *Sur un theoreme de Gluskin-Meyer-Pajor*, *C. R. Acad. Sci. Paris. Ser. I Math.* **317** (1993), pp. 155–158.
- [5] J. J. Schäffer, *Norms and determinants of linear mappings*, *Math. Z.*, **118** (1970), pp. 331–339.
- [6] I. V. Videnskii, N. A. Shirokov, *An extremal problem in the Wiener algebra*, *St. Petersburg Math. J.*, **11** (2000), No. 6, pp. 1035–1049.

Address:

Alexander Volberg
 Department of Mathematics
 Michigan State University
 East Lansing, Michigan 48824, USA
 volberg@math.msu.edu